

# *B* meson form factors from HQET simulations\*

C. Bernard,<sup>a</sup> C. DeTar,<sup>b</sup> Steven Gottlieb,<sup>c</sup> U. M. Heller,<sup>d</sup> J. Hetrick,<sup>a</sup> B. Jegerlehner,<sup>c</sup> C. McNeile,<sup>b</sup> K. Rummukainen,<sup>e</sup> R. Sugar,<sup>f</sup> D. Toussaint<sup>g</sup> and M. Wingate<sup>h</sup>

<sup>a</sup>Department of Physics, Washington University, St. Louis, MO 63130, USA

<sup>b</sup>Physics Department, University of Utah, Salt Lake City, UT 84112, USA  
and Zentrum für interdisziplinäre Forschung, Universität Bielefeld, Bielefeld, Germany

<sup>c</sup>Department of Physics, Indiana University, Bloomington, IN 47405, USA

<sup>d</sup>SCRI, The Florida State University, Tallahassee, FL 32306-4130, USA

<sup>e</sup>Fakultät für Physik, Universität Bielefeld, Bielefeld, Germany

<sup>f</sup>Department of Physics, University of California, Santa Barbara, CA 93106, USA

<sup>g</sup>Department of Physics, University of Arizona, Tucson, AZ 85721, USA

<sup>h</sup>Physics Department, University of Colorado, Boulder, CO 80309, USA

We use simulations of heavy quark effective field theory to calculate the Isgur-Wise function, and we demonstrate the feasibility of calculating the matrix element for the  $B \rightarrow \pi + l^+ \nu_l$  decay in the lattice heavy quark effective theory (HQET).

## 1. INTRODUCTION

We describe the calculation of the hadronic matrix elements that are required for the extraction of the  $V_{cb}$  and  $V_{ub}$  CKM matrix elements from experimental data [1]. To reach the bottom quark mass our strategy is to interpolate between results from relativistic quarks with  $m_q \leq m_c$  and results from lattice HQET [2]. Here we discuss only the HQET simulations, as our clover form factor simulations have only just started.

All of our simulations use  $n_f = 2$  dynamical staggered configurations with a volume  $16^3 \times 48$  and  $\beta = 5.445$ .

## 2. ISGUR-WISE FUNCTION

The Isgur-Wise function is the QCD matrix element required in the extraction of  $V_{cb}$  from experimental data. Experimental measurements of the slope of the Isgur-Wise function vary from 0.31 to

1.17, and the variations in theoretical predictions are nearly as large [3]. Initial attempts to calculate the Isgur-Wise function in lattice HQET had problems either with the signal to noise ratio [4] or the renormalization factors [7]. The first complete calculation has been done recently by the Kentucky group [8].

We use the same method as the Kentucky group (see also [4,7]). We ran at all permutations of the following velocities:  $(0,0,0)$ ,  $(0.1,0,0)$ ,  $(0.25,0,0)$  and  $(0.5,0,0)$ . Our sample size is 80 configurations, and our Wilson  $\kappa$  values are 0.160 and 0.163. A relative smearing function of  $e^{-0.67r}$  was used between the quarks in the  $B$  meson. In Fig. 1 we plot the bare Isgur-Wise function for various time separations between the current and the  $B$  source. If the ground state has been isolated, then the Isgur-Wise function should be independent of this separation. The data for  $\Delta t = 2, 3, 4$  are consistent within present errors.

It is traditional to report the slope of the Isgur-Wise function as a function of the dot product of

---

\*Presented by C. McNeile.




Figure 1. Unrenormalized Isgur-Wise function

$\Delta t$	$\kappa = 0.160$		$\kappa = 0.163$	
	$\rho^2$	$\chi^2/df$	$\rho^2$	$\chi^2/df$
2	0.415(8)	200/44	0.412(9)	171/44
3	0.48(4)	31/44	0.47(4)	24/44
4	0.48(12)	18/44	0.48(13)	17/44

Table 1

Preliminary fits to bare Isgur-Wise function data

the two meson velocities ( $\omega = v \cdot u$ ). In Table 1 we report fits to

$$\xi(\omega) = 1 - \rho^2(\omega - 1) \quad (1)$$

Assuming negligible quark mass dependence, our best estimate is therefore  $\rho^2 = 0.48(13)$  at the physical light quark mass. (We stress that it can not be compared with the experimental value until the renormalization factors calculated in [8] are included.) For comparison, at  $\beta = 6.0$  the Kentucky group gets a bare  $\rho^2 = 0.56$  and Hashimoto and Matsufuru [7] get  $\rho^2 \sim 0.54$  (where we have approximately removed the effect of tadpole improvement from their result [8]).

All the simulations of lattice HQET [7,8] show a very weak dependence of the slope on the light quark mass. However, the UKQCD collaboration [9] found a statistically significant decrease in  $\rho^2$  with light quark mass in their simulations

Vel	NP	$PT_{bare}$	$PT_{boost}$	tadpole
0.1	0.04(2)	0.074	0.051	0.085
0.25	0.13(2)	0.18	0.12	0.21
0.5	0.25(3)	0.34	0.23	0.41

Table 2

Various estimates of the HQET velocity renormalization

that used clover quarks for the  $b$  quark. We also tried fitting our bare Isgur-Wise function data to a fit model that had quadratic corrections of  $\omega$  in Eq. 1. Acceptable fits were found with approximately the same slope as in Table 1 and positive curvature.

The velocity of the lattice HQET action is renormalized [10] because the action breaks Lorentz symmetry. As we described last year [11], we have tried to estimate the renormalized velocity from the dispersion relation of an HQET meson at finite residual momentum [7]. The renormalized velocity can be implicitly defined from

$$E(\underline{p}, v^R) - E(0, v^R) = \frac{v^R \cdot \underline{p}}{v_0^R} \quad (2)$$

where  $E(\underline{p}, v^R)$  is the energy of the HQET meson at finite residual momentum ( $\underline{p}$ ). In Table 2 we show the results of the non-perturbative velocity renormalization. We fit all the correlators in the time region 5 to 13 and obtained correlated  $\chi^2/df$  slightly less than one. For comparison, we also show the results for the perturbative renormalization calculated by Mandula and Ogilvie [5,6], using both a boosted ( $g^2/u_0^4$ )  $PT_{boost}$  and bare coupling  $PT_{bare}$ , as well as using the tree-level tadpole improved estimate [8].

The results in Table 2 show that the velocity renormalization is large. These results would suggest that perturbation theory with a boosted coupling agrees best with the non-perturbative result. However, other analyses found better agreement between the tree-level tadpole scheme [8] and the non-perturbative calculations [5–7]. This issue is under investigation.

### 3. $B \rightarrow \pi$ FORM FACTOR

The observation of the decay  $B \rightarrow \pi + l^+ \nu_l$  allows a determination of  $V_{ub}$ , if the relevant QCD form factors can be calculated. There have been a number of lattice QCD calculations of the required form factors (see [1,12] for reviews). However, previous approaches suffer from the drawback that calculations are done at large  $q^2$ , thus requiring a large extrapolation to  $q^2 \approx 0$ , where measurements are currently made. To reach a low  $q^2$  requires large meson velocities—not easily achieved for heavy mesons in the NRQCD- or propagating-quark-approaches. As we have shown [11], a good signal can be obtained for the HQET  $B$  meson with a large velocity ( $v \approx 0.8$ ), so we propose the use of HQET—light simulations to explore lower  $q^2$  (see [13] for similar ideas). It is not clear that HQET will be a good approximation to the dynamics of the  $B$  meson, at these values of  $q^2$ , nor that a sufficiently good signal will be obtained. However, results should be very useful in helping to reduce the heavy quark extrapolation errors over simulations that only use clover quarks.

Because  $B \rightarrow$  light meson form factors have never been studied before using lattice HQET (although the static limit was studied in [14]), we have computed the matrix element for  $B \rightarrow \pi + l^+ \nu_l$  using HQET to check for a signal. We use the setup described in [14] with the heavy clover quark replaced by a HQET quark. In Fig. 2 we plot the ratio of three point functions to two point functions

$$\frac{\langle C_3(t; t_f) \rangle}{\langle C_2(t)_L \rangle \langle C_2(t_f - t)_B \rangle} \quad (3)$$

that is proportional to the  $\langle B | J_\mu | \pi \rangle$  matrix element, as a function of the operator time  $t$ . The  $B$  source is fixed at  $t_f = 23$ , and the light meson source is fixed at  $t = 0$ .

This work is supported by the DOE and the NSF. The computations were carried out at CCS (ORNL). This presentation was prepared with the kind additional support and assistance of the University of Bielefeld and its Zentrum für Interdisziplinäre Forschung.




Figure 2. HQET—light matrix element

### REFERENCES

1. T. Onogi, these proceedings.
2. J. Mandula and M. Ogilvie, *Phys. Rev. D* **45** (1992) 2183.
3. M. Neubert (1997) hep-ph/9702375.
4. J. Mandula and M. Ogilvie, *Nucl. Phys. B (Proc. Suppl.)* **34** (1994) 480.
5. J. E. Mandula and M. C. Ogilvie, (1996) hep-lat/9602004.
6. J. E. Mandula and M. C. Ogilvie, (1997) hep-lat/9703020,
7. S. Hashimoto and H. Matsufuru, *Phys. Rev. D* **54** (1996) 4578.
8. T. Draper and C. McNeile, (*Nucl. Phys. B (Proc. Suppl.)* **47** (1996) 429. J. Christensen, T. Draper and C. McNeile, these proceedings.
9. K. C. Bowler (UKQCD collaboration), *Phys. Rev. D* **52** (1995) 5067.
10. U. Aglietti, *Nucl. Phys.* **B421** (1994) 191.
11. C. Bernard *et al.*, *Nucl. Phys. B (Proc. Suppl.)* **53**, (1997) 374.
12. J. Flynn, *Nucl. Phys. B (Proc. Suppl.)* **53** (1997) 168.
13. J. Sloan, these proceedings.
14. UKQCD Collaboration, *Nucl. Phys.* **425** (1995) 4905.